



# SPENCERPORT CENTRAL SCHOOL DISTRICT

2019-2020 School Year

## New York State Immunization Requirements for School Entrance/Attendance

Children in a prekindergarten setting should be age-appropriately immunized. The number of doses depends on the schedule recommended by the Advisory Committee on Immunization Practices (ACIP). For grades pre-k through 11, intervals between doses of vaccine should be in accordance with the ACIP-recommended immunization schedule for persons 0 through 18 years of age. Doses received before the minimum age or intervals are not valid and do not count toward the number of doses listed below. Intervals between doses of vaccine DO NOT need to be reviewed for grade 12 except for interval between measles vaccine doses.

<b>Prekindergarten (Day Care, Head Start, Nursery or Pre-K)</b>	
Diphtheria and Tetanus toxoid-containing vaccine and Pertussis Vaccine (DTaP/DTP)	<b>4 doses</b>
Polio vaccine (IPV/OPV)	<b>3 doses</b>
Measles, Mumps and Rubella vaccine (MMR)	<b>1 dose</b>
Hepatitis B vaccine	<b>3 doses</b>
Varicella (Chickenpox) vaccine	<b>1 dose</b>
Hemophilus influenzae type b conjugate vaccine (Hib)	<b>1 to 4 doses</b>
Pneumococcal Vaccine (PCV)	<b>1 to 4 doses</b>
<b>Kindergarten through Grade 2</b>	
Diphtheria and Tetanus toxoid-containing vaccine and Pertussis Vaccine (DTaP/DTP)	<b>5 doses or 4 doses</b> if the 4 <sup>th</sup> dose was received at 4 years of age or <b>3 doses</b> if series started at 7 years of age or older
Polio vaccine (IPV/OPV)	<b>4 doses or 3 doses</b> if the 3 <sup>rd</sup> dose was received at 4 years of age or older
Measles, Mumps and Rubella vaccine (MMR)	<b>2 doses</b> (2 measles, 2 mumps 1 rubella)
Hepatitis B vaccine	<b>3 doses</b>
Varicella (Chickenpox) vaccine	<b>2 doses</b>
<b>Grades 3 through 5</b>	
Diphtheria and Tetanus toxoid-containing vaccine and Pertussis Vaccine (DTaP/DTP)	<b>5 doses or 4 doses</b> if the 4 <sup>th</sup> dose was received at 4 years of age or <b>3 doses</b> if series started at 7 years of age or older
Polio vaccine (IPV/OPV)	<b>3 doses</b>
Measles, Mumps and Rubella vaccine (MMR)	<b>2 doses</b> (2 measles, 2 mumps 1 rubella)
Hepatitis B vaccine	<b>3 doses</b>
Varicella (Chickenpox) vaccine	<b>1 dose</b>
<b>Grades 6 through 8</b>	
Diphtheria and Tetanus toxoid-containing vaccine and Pertussis Vaccine (DTaP/DTP)	<b>3 doses</b>
Tetanus and Diphtheria toxoid-containing vaccine and Pertussis vaccine booster (Tdap)	<b>1 dose</b>
Polio vaccine (IPV/OPV)	<b>4 doses or 3 doses</b> if the 3 <sup>rd</sup> dose was received at 4 years of age or older
Measles, Mumps and Rubella vaccine (MMR)	<b>2 doses</b> (2 measles, 2 mumps 1 rubella)
Hepatitis B vaccine	<b>3 doses or 2 doses</b> of adult hepatitis B vaccine (Recombivax) for children who received the doses at least 4 months apart between the ages of 11 - 15 years of age
Varicella (Chickenpox) vaccine	<b>2 doses</b>
<b>Grades 7 and 12</b>	
Meningococcal vaccine	<b>1 dose</b> in 7 <sup>th</sup> grade <b>2 doses</b> in 12 <sup>th</sup> grade one being a booster or <b>1 dose</b> on or after the age of 16
<b>Grades 9 through 12</b>	
Diphtheria and Tetanus toxoid-containing vaccine and Pertussis Vaccine (DTaP/DTP)	<b>3 doses</b>
Tetanus and Diphtheria toxoid-containing vaccine and Pertussis vaccine booster (Tdap)	<b>1 dose</b>
Polio vaccine (IPV/OPV)	<b>3 doses</b>
Measles, Mumps and Rubella vaccine (MMR)	<b>2 doses</b> (2 measles, 1 mumps 1 rubella)
Hepatitis B vaccine	<b>3 doses or 2 doses</b> of adult hepatitis B vaccine (Recombivax) for children who received the doses at least 4 months apart between the ages of 11 - 15 years of age
Varicella (Chickenpox) vaccine	<b>1 dose</b>

*Our Mission is to educate and inspire each student to love learning, pursue excellence and use knowledge, skills and attitudes to contribute respectfully and confidently to an ever-changing global community.*